



Queensland University of Technology
Brisbane Australia

This is the author's version of a work that was submitted/accepted for publication in the following source:

Keenan, Bethany E., Izatt, Maree T., Askin, Geoffrey N., Labrom, Robert D., Percy, Mark J., & Adam, Clayton J.

(2014)

The effect of endplate preselection when measuring supine versus standing cobb angle change in idiopathic scoliosis. In

Raso, Jim & Aubin, Carl-Eric (Eds.)

Proceedings of the 10th Meeting of the International Research Society of Spinal Deformities (IRSSD 2014), Hokkaido University Press, Sapporo, Japan.

This file was downloaded from: <http://eprints.qut.edu.au/78078/>

© Copyright 2015 Keenan et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Notice: *Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source:*

<http://dx.doi.org/10.1186/1748-7161-10-S1-O43>

THE EFFECT OF ENDPLATE PRESELECTION WHEN MEASURING SUPINE VERSUS STANDING COBB ANGLE CHANGE IN IDIOPATHIC SCOLIOSIS

*Keenan BE, Izatt MT, Askin GN, Labrom RD, Pearcy MJ, Adam CJ

QUT/Mater Paediatric Spine Research Group, Institute of Health and Biomedical Innovation, Queensland University of Technology and Mater Health Services, Brisbane, QLD, Australia.

Objectives

The primary aim of this study was to determine whether endplate pre-selection makes a difference to the Cobb Angle change between supine and standing which is known to occur in idiopathic scoliosis. A secondary aim of this study was to identify which (if any) patient characteristics were correlated with supine *versus* standing Cobb change.

Methods

Female Adolescent Idiopathic Scoliosis (AIS) patients with right-sided thoracic major curves were included in the retrospective study. Clinically measured Cobb Angles from existing standing coronal radiographs and fulcrum bending radiographs [1] were compared to existing low-dose supine CT scans taken within 3 months of the reference radiograph. Reformatted coronal CT images were used to measure supine Cobb Angle variability with and without endplate pre-selection (end-plates selected on the radiographs used on the CT images). Inter and intra-observer measurement variability was assessed. Multi-linear regression was used to investigate whether there was a relationship between supine to standing Cobb Angle change and patient characteristics (SPSS, v.21, IBM, USA).

Results

Fifty-two patients were included, with mean age of 14.6 (SD 1.8) years; all curves were Lenke Type 1 with mean Cobb Angle on supine CT of 41.8° (SD 6.4°) and 51.9° (SD 6.7°) on standing radiographs. The mean fulcrum bending Cobb Angle for the group was 22.6° (SD 7.5°). The 10° increase from supine to standing is consistent with existing literature. Pre-selecting vertebral endplates was found to increase the mean signed Cobb change by 0.6° (SD 2.3, range -9 to 6)° and the absolute Cobb change by 1.4° (SD 1.9, range 0 to 9)°. Multi-linear regression revealed a statistically significant relationship between supine to standing Cobb Angle change with: fulcrum flexibility ($p=0.001$), age ($p=0.027$) and standing Cobb Angle ($p<0.001$). In patients with high fulcrum flexibility scores, the supine to standing Cobb Angle change was as great as 20°. The 95% confidence intervals for intra-observer and inter-observer measurement variability were 3.1° and 3.6°, respectively.

Conclusion

Pre-selecting vertebral endplates causes only has a minor effect on supine to standing Cobb change in scoliosis. There is a statistically significant relationship between supine to standing Cobb Angle change and fulcrum flexibility. Therefore, supine to standing Cobb Angle change can be considered as a measure of spinal flexibility when both standing and supine images are clinically available.

References

1. 1997 Cheung KM, Luk KD. *JBJS (Am)*, 79; 1144-50